POSTPIL®

Effective and convenient 1-pill solution for emergency contraceptive

EASY-TO-USE · EASY-TO-FIND · EFFECTIVE · SAFE



WHAT IS POSTPIL®

- **Emergency contraceptive** to use after unprotected intercourse or in case of known or suspected contraceptive failure
- Package contains 1 tablet of levonorgestrel 1.5 mg
- Not an abortion pill; does not terminate established pregnancy
- Does not protect against sexually transmitted infections (STIs)
- Debunking common myths:
 - No health risks with repeated uses
 - Does not affect fertility
 - Does not increase the risk of ectopic pregnancy, cancer or cause long term side effects

WHO CAN USE POSTPIL®

- Women of all reproductive ages who want to avoid pregnancy, including adolescents, women who are breastfeeding, or women with history of ectopic pregnancy
- No absolute contraindications, except hypersensitivity to levonorgestrel

WHY USE POSTPIL®

- WHO-recommended regimen to prevent pregnancy
- Meets European standards and certifications for quality control of medicines and production processes
- 1-pill formulation is easier to follow correctly than 2-pill formulation
- High-quality yet affordable product

ALSO AVAILABLE FROM WOMANCARE GLOBAL



MISOMIFE-FEM® COMBO

Combi pack of 1 mifepristone 200 mg tablet and 4 misoprostol 200 mcg tablets Product Code: COMBO



INJECTA-FEM

Intramuscular contraceptive injection Medroxyprogesterone acete 150 mg/ml Product Code: INJECTA



IPAS DOUBLE VALVE ASPIRATOR

For uterine evacuation Product Code: DVS-10 or DVS-SU



POSTPIL®

Effective and convenient 1-pill solution for emergency contraceptive

EASY-TO-USE · EASY-TO-FIND · EFFECTIVE · SAFE



WHAT IS POSTPIL®

- **Emergency contraceptive** to use after unprotected intercourse or in case of known or suspected contraceptive failure
- Package contains 1 tablet of levonorgestrel 1.5 mg
- Not an abortion pill; does not terminate established pregnancy
- Does not protect against sexually transmitted infections (STIs)
- Debunking common myths:
 - No health risks with repeated uses
 - Does not affect fertility
 - Does not increase the risk of ectopic pregnancy, cancer or cause long term side effects

WHO CAN USE POSTPIL®

- Women of all reproductive ages who want to avoid pregnancy, including adolescents, women who are breastfeeding, or women with history of ectopic pregnancy
- No absolute contraindications, except hypersensitivity to levonorgestrel

WHY USE POSTPIL®

- WHO-recommended regimen to prevent pregnancy
- **Meets European standards and certifications** for quality control of medicines and production processes
- 1-pill formulation is easier to follow correctly than 2-pill formulation
- High-quality yet affordable product

ALSO AVAILABLE FROM WOMANCARE GLOBAL



MISOMIFE-FEM® COMBO

Combi pack of 1 mifepristone 200 mg tablet and 4 misoprostol 200 mcg tablets Product Code: COMBO



INJECTA-FEM

Intramuscular contraceptive injection Medroxyprogesterone acete 150 mg/ml Product Code: INJECTA



IPAS DOUBLE VALVE ASPIRATOR

For uterine evacuation Product Code: DVS-10 or DVS-SU



POSTPIL®

Effective and convenient 1-pill solution for emergency contraceptive

EASY-TO-USE · EASY-TO-FIND · EFFECTIVE · SAFE



WHEN TO USE POSTPIL®

- Can be used at any time during the menstrual cycle for the following situations:
 - Desire to avoid pregnancy
 - Nonconsensual or forced unprotected sex
 - Possible contraceptive failure or incorrect use
 - Concern about the withdrawal method
 - Uncertainty about the 'safe period'
- Can be used if already administered in the same menstrual cycle

HOW POSTPIL® WORKS

- Disrupts ovulation and thickens the cervical mucus, which inhibits sperm mobility and prevents fertilization
- No examinations or laboratory tests needed¹
- Reduces risk of pregnancy by 80%-90%
- **Side effects may include**: nausea, lower abdominal pain, fatigue, headache, dizziness, breast tenderness, delay of menses (>7 days)

HOW TO USE POSTPIL®

- To be used as soon as possible up to 3 days (72 hours) after unprotected intercourse for best results. Postpil® is more effective the sooner it is taken after unprotected sex.
- Dosage should not be taken more than once within 24 hours, except if the woman vomits within 2 hours after use
- Risk of pregnancy if repeated unprotected intercourse within 24 hours after taking the Postpil®

¹ Rodrigues, Isabel & Grou, Fabienne & Joly, Jacques. (2001). Effectiveness of emergency contraceptive pills between 72 and 120 hours after unprotected sexual intercourse. American journal of obstetrics and gynecology. 184. 531-7. 10.1067/mob.2001.111102.1

WomanCare Global Trading CIC is a UK Registered Community Interest Company (No.07231920). Email: contact@dktwomancare.org www.dktwomancare.org ALSO AVAILABLE FROM WOMANCARE GLOBAL



LEVOPLANT™

WHO-prequalified 2-rod hormonal implant Product Code: LEVO



MISO-FEM®

For uterine evacuation Pack of 4 misoprostol 200 mcg tablets Product Code: MISO



IPAS MVA PLUS® ASPIRATOR

Uterine evacuation device Product Code: PLUS

